

**Motion Problems**

Read from **Lesson 6** of the **1-D Kinematics** chapter at **The Physics Classroom**:

<http://www.physicsclassroom.com/Class/1DKin/U1L6a.html>

<http://www.physicsclassroom.com/Class/1DKin/U1L6b.html>

<http://www.physicsclassroom.com/Class/1DKin/U1L6c.html>

<http://www.physicsclassroom.com/Class/1DKin/U1L6d.html>

**MOP Connection:**      None

**Show your work on the following problems.**

1. An airplane accelerates down a runway at  $3.20 \text{ m/s}^2$  for 32.8 s until it finally lifts off the ground. Determine the distance traveled before take-off.
  
  
  
  
  
  
  
  
  
  
2. A race car accelerates uniformly from  $18.5 \text{ m/s}$  to  $46.1 \text{ m/s}$  in 2.47 seconds. Determine the acceleration of the car and the distance traveled.
  
  
  
  
  
  
  
  
  
  
3. A feather is dropped on the moon from a height of 1.40 meters. The acceleration of gravity on the moon is  $1.67 \text{ m/s}^2$ . Determine the time for the feather to fall to the surface of the moon.
  
  
  
  
  
  
  
  
  
  
4. A bullet leaves a rifle with a muzzle velocity of  $521 \text{ m/s}$ . While accelerating through the barrel of the rifle, the bullet moves a distance of 0.840 m. Determine the acceleration of the bullet (assume a uniform acceleration).

## 1-D Kinematics

- An engineer is designing a runway for an airport. Several planes will use the runway and the engineer must design it so that it is long enough for the largest planes to become airborne before the runway ends. If the largest plane accelerates at  $3.30 \text{ m/s}^2$  and has a takeoff speed of  $88.0 \text{ m/s}$ , then what is the minimum allowed length for the runway?
- A student drives 4.8-km trip to school and averages a speed of  $22.6 \text{ m/s}$ . On the return trip home, the student travels with an average speed of  $16.8 \text{ m/s}$  over the same distance. What is the average speed (in  $\text{m/s}$ ) of the student for the two-way trip? (Be careful.)
- Renatta Gas is driving through town at  $25.0 \text{ m/s}$  and begins to accelerate at a constant rate of  $-1.0 \text{ m/s}^2$ . Eventually Renatta comes to a complete stop. Represent Renatta's accelerated motion by sketching a velocity-time graph. Use kinematic equations to calculate the distance which Renatta travels while decelerating. Then use the velocity-time graph to determine this distance. **PSYW**
- Otto Emissions is driving his car at  $25.0 \text{ m/s}$ . Otto accelerates at  $2.0 \text{ m/s}^2$  for 5 seconds. Otto then maintains a constant velocity for 10 more seconds. Determine the distance Otto traveled during the entire 15 seconds. (Consider using a velocity-time graph.)
- Chuck Wagon travels with a constant velocity of  $0.5 \text{ mile/minute}$  for 10 minutes. Chuck then decelerates at  $-0.25 \text{ mile/min}^2$  for 2 minutes. Determine the total distance traveled by Chuck Wagon during the 12 minutes of motion. (Consider using a velocity-time graph.)